



Temperature Change Quantiles

20% quantile of temperature change [°C]
(2080-2100 vs 1980-2000)

DJF




JJA





Exceedance Probabilities

Probability of exceeding 2°C temperature change
(2080-2100 vs 1980-2000)

DJF





JJA





0.1 0.2 0.3 0.4 0.5 0.6 0.7 0.8 0.9


Exceedance Probabilities




Exceedance Fractions



Regional Assessment



Global Assessment



Source: AR4, IPCC

Model Extensions

- Use “more” data
 - ~> ensemble runs, model bias and internal variability, model present and future individually, . . .
- Use AOGCM specific weighting
 - ~> performance, “core” similarities, . . .
- Parameterize covariance matrices
 - ~> built in range, nonstationarity, . . .
- Building bi-/multivariate models
 - ~> use temperature for precipitation prediction, . . .
- Address computational complexity
 - ~> sparsity, GMRF, Metropolis-Hastings steps, . . .

References

Furrer, Knutti, Sain, Nychka, Meehl, (2007). Spatial patterns of probabilistic temperature change projections from a multivariate Bayesian analysis, *Geophys. Res. Lett.*, 34, L06711, doi:10.1029/2006GL027754.

Furrer, Sain, Nychka, Meehl, (2007). Multivariate Bayesian Analysis of Atmosphere-Ocean General Circulation Models, to appear in *Environmental and Ecological Statistics*.

Furrer, Sain, (2007). Spatial Model Fitting for Large Datasets with Applications to Climate and Microarray Problems, submitted to *Statistics and Computing*.

Sain, Furrer, Cressie, (2007). Combining Regional Climate Model Output via a Multivariate Markov Random Field Model. 56th Session of the International Statistical Institute, Lisbon, Portugal.